

Interaction Report

Report ID:
Date Produced: 02 November 2024

Antiretroviral Treatment**Co-medications**

Atazanavir alone (ATV)

Omeprazole

This report lists the summaries of potential interactions (i.e. "red", "amber" and "yellow" classifications) for the drugs in the table above.

Interactions with a "green" or "grey" classification (i.e. no clinically significant interaction or no clear data) have been checked and are listed at the end of this report, but summaries are not shown.

For full details of all interactions, see www.hiv-druginteractions.org.

Description of the interactions**Drugs that should not be coadministered (RED)****Atazanavir alone (ATV) + Omeprazole**

Coadministration of atazanavir with proton pump inhibitors is not recommended. Administration of omeprazole (40 mg once daily) decreased atazanavir C_{max}, AUC and C_{min} by 96%, 94% and 95%, respectively when given 2 h before atazanavir (400 mg, once daily). Omeprazole C_{max} and AUC increased by 24% and 45% when omeprazole (40 mg single dose) was given 2 hours after atazanavir (400 mg once daily). Coadministration is not recommended in treatment experienced patients. If coadministration is judged unavoidable in treatment-naive patients, close clinical monitoring is recommended and doses of omeprazole should not exceed 20 mg and must be taken approximately 12 hours prior to the atazanavir.