

Interaction Report

Report ID:

Date Produced:

19 April 2025

Antiretroviral Treatment

Co-medications

Efavirenz (EFV)

Furosemide

This report lists the summaries of potential interactions (i.e. "red", "amber" and "yellow" classifications) for the drugs in the table above.

Interactions with a "green" or "grey" classification (i.e. no clinically significant interaction or no clear data) have been checked and are listed at the end of this report, but summaries are not shown.

For full details of all interactions, see www.hiv-druginteractions.org.

Description of the interactions

Potential weak interaction - additional action/monitoring or dosage adjustment is unlikely to be required (YELLOW)

Efavirenz (EFV) + Furosemide

Coadministration has not been studied but based on metabolism and clearance a pharmacokinetic interaction is unlikely. Furosemide is glucuronidated mainly in the kidney (UGT1A9) and to a lesser extent in the liver (UGT1A1). A large proportion of furosemide is also eliminated unchanged really. Furosemide has a conditional risk of QT prolongation and/or TdP on the CredibleMeds.org website. Efavirenz was shown to prolong the QT interval above the regulatory threshold of concern in homozygous carriers of the CYP2B6*6/*6 allele (i.e. 516T variant in the gene encoding CYP2B6). The European product label for efavirenz contraindicates coadministration with a drug with a known risk of Torsade de Pointes whereas the American product label for efavirenz recommends that alternatives should be considered. As the potential risk of a QT interval prolongation relates specifically to homozygous carriers of CYP2B6*6/*6 and given the accumulated years of safety data with efavirenz and such drugs, the contraindication is not reflected in the colour coding of this interaction summary.